



ІНСТИТУТ  
ФІЗИКИ  
КОНДЕНСОВАНИХ  
СИСТЕМ

ICMP-05-15E

A. Duviryak

Large- $j$  expansion method for two-body Dirac equation

УДК: 531/533; 530.145; 531.18; 539.12

РАС: 03.65.Pm, 03.65.Ge, 12.39.Pn

Метод розкладів за великим  $j$  для двочастинкового рівняння Дірака

А. Дувіряк

**Анотація.** Двочастинкове рівняння Дірака в координатному представленні зведено, з врахуванням його симетрійних властивостей, до зв'язаної пари радіальних диференціальних рівнянь 2-го порядку. До цієї пари застосовано метод розкладів за великим  $j$  з метою розв'язання задачі зв'язаних станів.

**Large- $j$  expansion method for two-body Dirac equation**

A. Duviryak

**Abstract.** Using symmetry properties the two-body Dirac equation in coordinate representation is reduced to the coupled pair of radial second-order differential equations. Then the large- $j$  expansion technique is used to solve a bound state problem. Linear-plus-Coulomb potentials of different spin structure are examined in order to describe the asymptotic degeneracy and fine splitting of light meson spectra.

Подається в SIGMA  
Submitted to SIGMA



## 1. Introduction

Two-body Dirac equations (2BDE), i.e., the Breit equations [1] and its generalizations [2–10], are used frequently for the description of relativistic bound state problem, especially in nuclear [11, 12] and hadronic [13–16] physics. Apart two free-particle Dirac terms the 2BDE may include potentials which are local matrix-functions in the coordinate representation. This form provide an intuitive understanding of the interaction and may suggest a proper physical choice of the potential in phenomenological models.

But the 2BDE are pathological if certain interaction terms are not treated perturbationally. The set of radially reduced equations [5, 7, 13] may possess non-physical energy-dependent poles at finite distance  $r$  between particles [17, 18]. Correspondingly, an exact boundary-value problem becomes incorrect mathematically or improper for assumed physical treatment.

Here we consider a possibility to avoid pathological peculiarities of 2BDE using a pseudo-perturbative technique similar to  $1/N$  [19–21] or  $1/\ell$  expansions [22, 23]. These methods are applicable to the case of a strong coupling and are little affected by boundary peculiarities of the boundary-value problem.

In our case natural expansion parameter is  $1/j$ , where  $j$  is the conserved total angular momentum. After the radial reduction is performed the 2BDE takes the form of the set of eight coupled first-order differential equations [5, 7]. Using a chain of transformations we reduce it to the pair of coupled second-order equations and apply the  $1/j$  expansion technique. The method is applied to the potential model of meson based on the 2BDE.

## 2. 2-body Dirac equation and its radial reduction

In the centre-of-mass reference frame the 2BDE has the form:

$$\{h_1(\mathbf{p}) + h_2(-\mathbf{p}) + U(\mathbf{r}) - E\} F(\mathbf{r}) = 0, \quad (1)$$

where  $F(\mathbf{r})$  is a 16-component wave function,

$$h_a(\mathbf{p}) = \boldsymbol{\alpha}_a \cdot \mathbf{p} + m_a \beta_a \equiv -i \boldsymbol{\alpha}_a \cdot \boldsymbol{\nabla} + m_a \beta_a, \quad a = 1, 2, \quad (2)$$

are Dirac Hamiltonians of free fermions of mass  $m_a$ ,  $a = 1, 2$ , and  $U(\mathbf{r})$  is an interaction potential. If  $F(\mathbf{r})$  is presented in  $4 \times 4$ -matrix representation, the operators  $\boldsymbol{\alpha}_a$  and  $\beta_a$  act as follows:  $\boldsymbol{\alpha}_1 F = \boldsymbol{\alpha} F$ ,  $\boldsymbol{\alpha}_2 F = F \boldsymbol{\alpha}^T$  etc, where  $\boldsymbol{\alpha}$  and  $\beta$  are Dirac matrices.

The potential  $U(\mathbf{r})$  is the Hermitian matrix-function, it is invariant under rotation and space inversion transformations (so that the total Hamiltonian  $H = h_1 + h_2 + U$  is too). The general form of it is parametrized by 48 scalar function of  $r = |\mathbf{r}|$  [10]. Of physical meaning are potentials admitting field-theoretical interpretation of interaction. In particular, potentials reflecting a spin structure of vector and scalar relativistic interactions are used frequently in potential quark models of mesons. We will consider such a model in the Section 5 using few examples of scalar and vector potentials known in a literature. In the present section the structure of potential is not essential.

In order to apply a pseudoperturbative expansion method to the 2-body Dirac equation let us transform it to an appropriate form.

First of all we perform a radial reduction. Following the Ref. [5, 7] we put the wave eigenfunction  $F(\mathbf{r})$  of the total angular momentum  $j$  and the parity  $P$  in the  $2 \times 2$  block-matrix form:

$$F(\mathbf{r}) = \frac{1}{r} \begin{bmatrix} i s_1(r) \phi^A(\mathbf{n}) + i s_2(r) \phi^0(\mathbf{n}) & t_1(r) \phi^-(\mathbf{n}) + t_2(r) \phi^+(\mathbf{n}) \\ u_1(r) \phi^-(\mathbf{n}) + u_2(r) \phi^+(\mathbf{n}) & i v_1(r) \phi^A(\mathbf{n}) + i v_2(r) \phi^0(\mathbf{n}) \end{bmatrix} \quad (3)$$

for the parity  $P = (-)^{j \pm 1}$  states, and similar form for the parity  $P = (-)^j$  states but with superscripts interchanged as follows:  $(A, 0) \leftrightarrow (-, +)$ . Here  $\mathbf{n} = \mathbf{r}/r$ , the bispinor harmonics  $\phi^A(\mathbf{n})$  corresponds to a singlet state with a total spin  $s = 0$  and an orbital momentum  $\ell = j$ , and  $\phi^0(\mathbf{n})$ ,  $\phi^-(\mathbf{n})$ ,  $\phi^+(\mathbf{n})$  correspond to  $s = 1$  and  $\ell = j, j+1, j-1$ . Then for  $j > 0$  the eigenstate problem (1) reduces to the set of eight first-order differential equations with the functions  $s_1(r) \dots v_2(r)$  and the energy  $E$  to be found.

It is convenient to present this set in the following matrix form. Let us introduce the 8-dimensional vector-function:  $\mathbf{X}(r) = \{s_1(r), s_2(r), t_1(r), \dots, v_2(r)\}$  Then the set of radial equations reads:

$$\left\{ \mathbf{H}(j) \frac{d}{dr} + \mathbf{V}(r, E, j) \right\} \mathbf{X}(r) = E \mathbf{X}(r), \quad (4)$$

where the  $8 \times 8$  real matrices  $\mathbf{H}(j)$  and  $\mathbf{V}(r, E, j) = \mathbf{G}(j)/r + \mathbf{m} + \mathbf{U}(r, j) - E \mathbf{I}$  possesses properties  $\mathbf{H}^T = -\mathbf{H}$ ,  $\mathbf{V}^T = \mathbf{V}$ , the diagonal matrix  $\mathbf{m} = \text{diag}(m_+ \mathcal{J}, m_- \mathcal{J}, -m_- \mathcal{J}, -m_+ \mathcal{J})$  (here  $\mathcal{J}$  is  $2 \times 2$  unit matrix and  $m_{\pm} = m_1 \pm m_2$ ) and  $j$ - and  $P$ -dependent matrices  $\mathbf{H}(j)$ ,  $\mathbf{G}(j)$  are constant (i.e., free of  $r$ ), and matrix-potential  $\mathbf{U}(r, j)$  comes from interacting term of the equation (1). For the case  $j = 0$  components  $s_2 = t_2 = u_2 = v_2 = 0$  so that the dimension of the problem (4) reduces from 8 to 4.

It turns out that  $\text{rank } \mathbf{H} = 4$  (2 for  $j = 0$ ). In other words, only four equations of the set (4) are differential while remaining ones are

algebraic. They can be split by means of some orthogonal (i.e., of  $O(8)$  group) transformation. In new basis we have

$$\mathbf{X}(r) = \begin{bmatrix} \mathbf{X}_1 \\ \mathbf{X}_2 \end{bmatrix}, \quad \mathbf{H} = 2 \begin{bmatrix} \mathbf{J}^{(2)} & \mathbf{0} \\ \mathbf{0} & \mathbf{0} \end{bmatrix}, \quad (5)$$

where  $\mathbf{J}^{(2)}$  is the symplectic  $4 \times 4$  matrix. Thus we arrive at the set

$$2\mathbf{J}^{(2)}\mathbf{X}'_1 + \mathbf{V}_{11}\mathbf{X}_1 + \mathbf{V}_{12}\mathbf{X}_2 = 0, \quad (6)$$

$$\mathbf{V}_{21}\mathbf{X}_1 + \mathbf{V}_{22}\mathbf{X}_2 = 0. \quad (7)$$

Eliminating  $\mathbf{X}_2$  from (6) by means of (7) we get a differential set for the 4-vector  $\mathbf{X}_1$

$$\left\{ \mathbf{J}^{(2)} \frac{d}{dr} + \mathbf{V}^\perp(r, E, j) \right\} \mathbf{X}_1(r) = 0, \quad \mathbf{V}^\perp = (\mathbf{V}_{11} - \mathbf{V}_{12}\mathbf{V}_{22}^{-1}\mathbf{V}_{21})/2 \quad (8)$$

while  $\mathbf{X}_2$  then follows from the algebraic relation  $\mathbf{X}_2 = -\mathbf{V}_{22}^{-1}\mathbf{V}_{21}\mathbf{X}_1$ .

The elimination of  $\mathbf{X}_2$  causes non-physical energy-dependent singular points (apart of  $r = 0$  and physical singularities of potentials) in matrix elements of  $\mathbf{V}^\perp$ .

Now we present the 4-vector  $\mathbf{X}_1$  in 2+2 block form,

$$\mathbf{X}_1(r) = \begin{bmatrix} \Phi_1 \\ \Phi_2 \end{bmatrix}, \quad \mathbf{V}^\perp = \begin{bmatrix} \mathcal{V}_{11} & \mathcal{V}_{12} \\ \mathcal{V}_{21} & \mathcal{V}_{22} \end{bmatrix}, \quad (9)$$

eliminate then  $\Phi_2$  and arrive at the second-order differential equations for 2-vector  $\Phi_1$ :

$$\mathcal{L}(E)\Phi_1 = \left\{ \left( \frac{d}{dr} + \mathcal{V}_{12} \right) [\mathcal{V}_{22}]^{-1} \left( \frac{d}{dr} - \mathcal{V}_{21} \right) + \mathcal{V}_{11} \right\} \Phi_1 = 0. \quad (10)$$

The matrix  $\mathcal{V}_{22}$  is diagonal for all potentials considered in Section 5 (and many other ones). In these cases we can perform the transformation:

$$\tilde{\Phi}_1 = \Phi_1 / \sqrt{\mathcal{V}_{22}}, \quad \tilde{\mathcal{L}} = \sqrt{\mathcal{V}_{22}} \mathcal{L} \sqrt{\mathcal{V}_{22}} \quad (11)$$

providing for the operator  $\tilde{\mathcal{L}}$  the form which is as close as possible to 2-term form:

$$\tilde{\mathcal{L}}(E) = \frac{d^2}{dr^2} - \mathcal{W}(r, E, j) - \{Z(r, E, j), \frac{d}{dr}\}_+ \mathcal{J}^{(1)}; \quad (12)$$

here  $\mathcal{W}(r, E, j)$  is a symmetric  $2 \times 2$  matrix,  $\mathcal{J}^{(1)}$  is  $2 \times 2$  symplectic matrix and  $\{\dots, \dots\}_+$  denotes the anticommutator.

We are nearly to apply the  $1/j$  expansion method to the equation (12). For physically interesting cases the function  $Z$  vanishes or it is negligible at  $j$  large. Thus the equation (12) has a  $2 \times 2$  matrix 2-term form. Before proceeding further we consider a simpler example of a single 2-term relativistic equation.

### 3. Todorov equation via $1/\ell$ method

Here we consider the Todorov-type equation describing the relativistic system of two interacting scalar particles in the centre-of-mass reference frame [24–26]:

$$\{\mathbf{p}^2 + U(r, E) - b(E)\} \Psi(\mathbf{r}) = 0. \quad (13)$$

Here  $\mathbf{p} = -i\nabla$ , the quasipotential  $U(r, E)$  depends on energy  $E$  of the system, and the binding parameter  $b(E)$  is the following function of  $E$ ,

$$b(E) = \frac{1}{4}E^2 - \frac{1}{2}(m_1^2 + m_2^2) + \frac{1}{4}(m_1^2 - m_2^2)^2/E^2, \quad \text{so that} \\ E(b) = \sum_{a=1}^2 \sqrt{m_a^2 + b}. \quad (14)$$

The corresponding radial equation takes the form

$$\left\{ \frac{d^2}{dr^2} - W(r, E, \ell) \right\} \psi(r) = 0, \quad (15)$$

where  $\ell$  is the angular momentum quantum number, and

$$W(r, E, \ell) = U(r, E) + \ell(\ell + 1)/r^2 - b(E). \quad (16)$$

Let us consider motion of the system in the neighbourhood of classical stable circular orbit. Given  $\ell > 0$ , the radius  $r_c = r_c(\ell)$  of the stable circular orbit and the corresponding energy  $E_c = E_c(\ell)$  satisfy conditions:

$$W(r_c, E_c, \ell) = 0, \quad \partial W(r_c, E_c, \ell)/\partial r = 0 \quad (17)$$

and  $\partial^2 W(r_c, E_c, \ell)/\partial r^2 > 0$ .

One puts  $r = r_c + \Delta r$  and  $E = E_c + \Delta E$  where  $\Delta r$  and  $\Delta E$  are small in some meaning, and expand the function  $W(r_c + \Delta r, E_c + \Delta E, \ell)$  in power series with respect  $\Delta r$  and  $\Delta E$ . Then due to the conditions (17) leading terms of this expansion represent the harmonic oscillator potential and other ones are anharmonic terms. If the conditions (17) hold for any large value of  $\ell$  it is possible by renormalization of  $\Delta r$  and  $\Delta E$  to single out

in the equation (15) the  $\ell$ -independent harmonic oscillator problem and anharmonic perturbations which disappear if  $\ell \rightarrow \infty$ . This is the idea of  $1/\ell$  expansion method. The application of pseudoperturbative techniques of this type [19–23] to our case meets two peculiarities: the equation (15) represents a nonlinear spectral problem, and an exact solution of the equations (17) may appear to be unknown or too cumbersome for practical use. Thus we modify the technique.

Let us introduce the parameter  $\lambda = 1/\sqrt{\ell}$  which is small at  $\ell$  large. Since the exact form of the functions  $r_c(\ell)$  and  $E_c(\ell)$  is unknown in general, we first determine asymptotics  $r_c \sim r_\infty(\lambda)$ ,  $b_c = b(E_c) \sim b_\infty(\lambda)$  at  $\lambda \rightarrow 0$  which may be found much easily. Then the functions  $r_c(\ell)$  and  $E_c(\ell)$  can be presented in the form:

$$r_c(\lambda) = r_\infty(\lambda)\rho(\lambda), \quad b_c(\lambda) = b_\infty(\lambda)\mu(\lambda), \quad (18)$$

$$\rho(\lambda) = 1 + \lambda\rho^{(1)} + \lambda^2\rho^{(2)} + \dots, \quad \mu(\lambda) = 1 + \lambda\mu^{(1)} + \lambda^2\mu^{(2)} + \dots \quad (19)$$

where expansion coefficients  $\rho^{(n)}$ ,  $\mu^{(n)}$ ,  $n = 1, 2, \dots$  (and thus the analytical functions  $\rho(\lambda)$  and  $\mu(\lambda)$ ) can be found, step by step, from the conditions:

$$\bar{W}(\rho, \mu, \lambda) = 0, \quad \partial\bar{W}(\rho, \mu, \lambda)/\partial\rho = 0 \quad (20)$$

and  $\partial^2\bar{W}(\rho, \mu, \lambda)/\partial\rho^2 > 0$ ; here the dimensionless function

$$\bar{W}(\rho, \mu, \lambda) = \lambda^4 r_\infty^2 W[r_\infty\rho, E(b_\infty\mu), 1/\lambda^2] \quad (21)$$

is regular at  $\lambda \rightarrow 0$  by construction.

Now we transit to the dimensionless variable  $r \rightarrow \xi$  and spectral parameter  $b(E) \rightarrow \epsilon$ ,

$$r = r_\infty(\lambda)[\rho(\lambda) + \lambda\xi], \quad b = b_\infty(\lambda)[\mu(\lambda) + \lambda^2\epsilon], \quad (22)$$

in terms of which the equation (15) takes the form

$$\left\{ \frac{d^2}{d\xi^2} - \frac{1}{\lambda^2} w(\xi, \epsilon, \lambda) \right\} \psi(\xi) = 0 \quad (23)$$

with

$$\psi(\xi) = \Psi[r_\infty(\rho + \lambda\xi)], \quad (24)$$

and

$$w(\xi, \epsilon, \lambda) = \bar{W}(\rho + \lambda\xi, \mu + \lambda^2\epsilon, \lambda). \quad (25)$$

If the functions  $\rho(\lambda)$  and  $\mu(\lambda)$  satisfy the conditions (20) the equation (23) is nonsingular at  $\lambda \rightarrow 0$ . This is true even if we use the first-order approximate solution to (20) in (22),

$$\rho(\lambda) = 1 + \lambda\rho^{(1)}, \quad \mu(\lambda) = 1 + \lambda\mu^{(1)}. \quad (26)$$

Indeed,

$$\begin{aligned} \frac{1}{\lambda^2} w(\xi, \epsilon, \lambda) &= \frac{1}{\lambda^2} \bar{W}[\rho(\lambda) + \lambda\xi, \mu(\lambda) + \lambda^2\epsilon, \lambda] \\ &= \frac{1}{\lambda^2} \bar{W}^{(0)} + \frac{1}{\lambda} \left\{ \frac{\partial\bar{W}^{(0)}}{\partial\rho}(\rho^{(1)} + \xi) + \frac{\partial\bar{W}^{(0)}}{\partial\mu}(\mu^{(1)} + \lambda\epsilon) + \frac{\partial\bar{W}^{(0)}}{\partial\lambda} \right\} + \\ &\quad + \frac{1}{2} \frac{\partial^2\bar{W}^{(0)}}{\partial\rho^2}(\rho^{(1)} + \xi)^2 + \frac{1}{2} \frac{\partial^2\bar{W}^{(0)}}{\partial\mu^2}[\mu^{(1)}]^2 + \frac{1}{2} \frac{\partial^2\bar{W}^{(0)}}{\partial\lambda^2} + \\ &\quad + \frac{\partial^2\bar{W}^{(0)}}{\partial\rho\partial\mu}(\rho^{(1)} + \xi)\mu^{(1)} + \frac{\partial^2\bar{W}^{(0)}}{\partial\rho\partial\lambda}(\rho^{(1)} + \xi) + \frac{\partial^2\bar{W}^{(0)}}{\partial\mu\partial\lambda}\mu^{(1)} + O(\lambda). \end{aligned} \quad (27)$$

Singular terms are absent if the following set of equations holds:

$$\bar{W}^{(0)} = 0, \quad \partial\bar{W}^{(0)}/\partial\rho = 0, \quad (28)$$

$$\frac{\partial\bar{W}^{(0)}}{\partial\mu}\mu^{(1)} + \frac{\partial\bar{W}^{(0)}}{\partial\lambda} = 0. \quad (29)$$

Besides, zero-order terms which are linear in  $\xi$  disappear if

$$\frac{\partial^2\bar{W}^{(0)}}{\partial\rho^2}\rho^{(1)} + \frac{\partial^2\bar{W}^{(0)}}{\partial\rho\partial\mu}\mu^{(1)} + \frac{\partial^2\bar{W}^{(0)}}{\partial\rho\partial\lambda} = 0. \quad (30)$$

Notice that the equations (28) and (29)–(30) represent the conditions (20) in the zeroth and first orders of  $\lambda$ , respectively. Thus the equations (28) hold identically and the (29)–(30) are linear equations with  $\rho^{(1)}$  and  $\mu^{(1)}$  to be found.

In zero-order approximation the equation (23) reduces to the harmonic oscillator problem

$$\left\{ \frac{d^2}{d\xi^2} + \kappa\epsilon - \nu - \omega^2\xi^2 \right\} \psi(\xi) = 0 \quad (31)$$

with

$$\kappa = -\frac{\partial\bar{W}^{(0)}}{\partial\mu}, \quad \omega^2 = \frac{1}{2} \frac{\partial^2\bar{W}^{(0)}}{\partial\rho^2}, \quad (32)$$

$$\begin{aligned} \nu &= -\frac{1}{2} \frac{\partial^2\bar{W}^{(0)}}{\partial\rho^2}[\rho^{(1)}]^2 + \frac{1}{2} \frac{\partial^2\bar{W}^{(0)}}{\partial\mu^2}[\mu^{(1)}]^2 \\ &\quad + \frac{1}{2} \frac{\partial^2\bar{W}^{(0)}}{\partial\lambda^2} + \frac{\partial^2\bar{W}^{(0)}}{\partial\mu\partial\lambda}\mu^{(1)}, \end{aligned} \quad (33)$$

$$\begin{aligned} \mu^{(1)} &= -\frac{\partial^2\bar{W}^{(0)}/\partial\lambda}{\partial^2\bar{W}^{(0)}/\partial\mu}, \\ \rho^{(1)} &= -\frac{1}{\partial^2\bar{W}^{(0)}/\partial\rho^2} \left\{ \frac{\partial^2\bar{W}^{(0)}}{\partial\rho\partial\mu}\mu^{(1)} + \frac{\partial^2\bar{W}^{(0)}}{\partial\rho\partial\lambda} \right\}, \end{aligned} \quad (34)$$

where  $\partial\bar{W}^{(0)}/\partial\mu = \lim_{\lambda\rightarrow 0} \partial\bar{W}/\partial\mu = \partial\bar{W}/\partial\mu(1, 1, 0)$  etc. The higher-order terms in the expansion (27) can be considered as perturbations to the oscillator problem (31). They depend, in general, on a spectral parameter  $\epsilon$  and can be taken into account by means of the perturbative procedure [25] which is appropriate to this case. Otherwise the treatment is closed to that of Refs. [19–23].

The eigenvalues in zero-order approximation  $\epsilon_{n_r} = [\omega(2n_r + 1) + \nu]/\kappa$ , where  $n_r = 0, 1, \dots$  is a radial quantum number, are to be corrected by means of higher orders of perturbative procedure. Then, using 2nd eq. of (22) in (14) gives us the energy spectrum.

#### 4. Breit-type equation via $1/j$ method

At this point we return to the radial 2BDE in the form  $\tilde{\mathcal{L}}(E)\tilde{\Phi}_1 = 0$ , where the  $2\times 2$  matrix operator  $\tilde{\mathcal{L}}(E)$  is given by Eq. (12) with the last term neglected. Let us put

$$\Phi_1 = \begin{bmatrix} \Psi_1 \\ \Psi_2 \end{bmatrix}, \quad (35)$$

where  $\Psi_1$  and  $\Psi_2$  are component of  $\Phi_1$ . Then the equation (10) can be presented as a pair of coupled ordinary second-order differential equations:

$$\frac{d^2}{dr^2}\Psi_1(r) - W_1(r, E, j)\Psi_1(r) = Y(r, E, j)\Psi_2(r), \quad (36)$$

$$\frac{d^2}{dr^2}\Psi_2(r) - W_2(r, E, j)\Psi_2(r) = Y(r, E, j)\Psi_1(r). \quad (37)$$

We will treat this system perturbationally using the pseudosmall parameter  $\lambda = 1/j$ .

Let us suppose for a moment that the right-hand side of the system (36)–(37) can be ignored, so that these equations decouple. Then we can apply to each of equations the scheme of the Section 3. We define radii and energies of circular orbits by means of the conditions:

$$W_i(r_i, E_i, j) = 0, \quad \frac{\partial W_i(r_i, E_i, j)}{\partial r} = 0, \quad \frac{\partial^2 W_i(r_i, E_i, j)}{\partial r^2} > 0, \quad (38)$$

$i = 1, 2$ . Then we single out asymptotics of these functions of  $\lambda$  by means of the relations:

$$r_i(\lambda) = r_{i\infty}(\lambda)\rho_i(\lambda), \quad b_i(\lambda) = b_{i\infty}(\lambda)\mu_i(\lambda), \quad (39)$$

$$\rho_i(\lambda) = 1 + \lambda\rho_i^{(1)} + \lambda^2\rho_i^{(2)} + \dots, \quad \mu_i(\lambda) = 1 + \lambda\mu_i^{(1)} + \lambda^2\mu_i^{(2)} + \dots, \quad (40)$$

and, using the relations

$$r = r_{i\infty}(\lambda)[\rho_i(\lambda) + \lambda\xi_i], \quad b = b_{i\infty}(\lambda)[\mu_i(\lambda) + \lambda^2\epsilon_i], \quad i = 1, 2 \quad (41)$$

we reformulate the equation (36) in terms of the dimensionless variable  $\xi_1$  and the spectral parameters  $\epsilon_1$  while the equation (37) – in terms of  $\xi_2$  and  $\epsilon_2$ . Finally, we perform an expansion of the equations in powers of  $\lambda$  and solve them separately.

Now we are going to take an actual coupling of the equations (36) and (37) into account. First of all we note that the variables  $\xi_1$  and  $\xi_2$  are not independent of one another, as well as the spectral parameters  $\epsilon_1$  and  $\epsilon_2$  are not. Thus we should choose common variables in both the equations.

Let us first choose  $\xi = \xi_1$ ,  $\epsilon = \epsilon_1$ . Then the set (36)–(37) reduces to the form:

$$\psi_1''(\xi) - \frac{1}{\lambda^2}w_1(\xi, \epsilon, \lambda)\psi_1(\xi) = y(\xi, \epsilon, \lambda)\psi_2(\xi), \quad (42)$$

$$\psi_2''(\xi) - \frac{1}{\lambda^2}w_2(\xi, \epsilon, \lambda)\psi_2(\xi) = y(\xi, \epsilon, \lambda)\psi_1(\xi), \quad (43)$$

where

$$\psi_i(\xi) = \Psi_i[r_{1\infty}(\rho_1 + \lambda\xi)], \quad i = 1, 2, \quad (44)$$

$$w_i(\xi, \epsilon, \lambda) = \lambda^4 r_{1\infty}^2 W_i[r_{1\infty}(\rho_1 + \lambda\xi), E(b_{1\infty}(\mu_1 + \lambda^2\epsilon)), 1/\lambda^2], \quad (45)$$

$$y(\xi, \epsilon, \lambda) = \lambda^2 r_{1\infty}^2 Y[r_{1\infty}(\rho_1 + \lambda\xi), E(b_{1\infty}(\mu_1 + \lambda^2\epsilon)), 1/\lambda^2]. \quad (46)$$

The functions (44)–(46) are regular at  $\lambda \rightarrow 0$ . Moreover, the general structure of the function  $w_1$  is the same as that of  $w$  in the Section 3 (see eqs. (25), (27)). In particular,  $w_1 = O(\lambda^2)$ . Thus the equation (42) is similar to (23) (but with non-zero right-hand side). It admits similar expansion in  $\lambda$ .

On the contrary, the function  $w_2$  may possess different behaviour at  $\lambda \rightarrow 0$ . Here we consider three cases.

1). Let  $r_{2\infty} \neq r_{1\infty}$  and  $b_{2\infty} \neq b_{1\infty}$ . Then  $w_2 = O(\lambda^{-n})$ ,  $n \geq 0$  (except perhaps very special examples which we do not consider). In this case one can solve formally the equation (43) in favour of  $\psi_2(\xi)$  as follows:

$$\psi_2 = - \left(1 - \frac{\lambda^2}{w_2} \frac{\partial^2}{\partial \xi^2}\right)^{-1} \frac{\lambda^2}{w_2} y \psi_1 = - \sum_{n=0}^{\infty} \left(\frac{\lambda^2}{w_2} \frac{\partial^2}{\partial \xi^2}\right)^n \frac{\lambda^2}{w_2} y \psi_1. \quad (47)$$

This representation leads to the loss of solutions for  $\psi_2$  which are not analytical in  $\lambda$  and thus have nothing to do with the perturbation procedure. The use of (47) in the r.-h.s. of (42) permits us to eliminate  $\psi_2$

from (42) and thus to obtain a close wave equation for  $\psi_1$ . The structure and treatment of this equation are the same as those of the equation (23). Moreover, it is obvious from (47) that at least the zero- and the first-order terms of  $\psi_2$  vanish. Thus the r.h.s. of (42) does not contribute in lower orders of perturbation procedure. In zero-order approximation we have the oscillator problem.

**2).** Let  $r_{2\infty} = r_{1\infty}$  and  $b_{2\infty} = b_{1\infty}$  but  $\rho_2 - \rho_1 = O(\lambda)$  and  $\mu_2 - \mu_1 = O(\lambda)$ . Then  $w_2 = O(\lambda)$ . Since  $\lambda^2/w_2 = O(\lambda)$  the perturbative treatment (47) of the equation (43) is still valid. The only difference from the case **1)** is that the r.h.s. of eq. (42) may contribute in the first order of  $\lambda$ .

In both the above cases we used the dimensionless variable  $\xi_1$  and obtained a closed eigenstate equation (which we will reference to as the problem **1)** for the wave function  $\psi_1(\xi_1)$  and the spectral parameter  $\epsilon_1$ . We can proceed with the variable  $\xi_2$  and obtain the problem **2)** for the function  $\psi_2(\xi_2)$  and the parameter  $\epsilon_2$ . One might be inclined to think that both the problem **1)** and **2)** are equivalent and lead to the same spectrum (in terms of energy  $E$ ). Actually, these problems complement one another. Indeed, it is evident from the relation:

$$\epsilon_2 - \epsilon_1 = \frac{1}{\lambda^2} \left\{ \frac{b_{1\infty}}{b_{2\infty}} \mu_1 - \mu_2 \right\} + \left\{ \frac{b_{1\infty}}{b_{2\infty}} - 1 \right\} \epsilon_1 \quad (48)$$

that in both **1)** and **2)** cases  $|\epsilon_2 - \epsilon_1| \rightarrow \infty$  if  $\lambda \rightarrow 0$ . It does mean that energy levels following from the problem **1)** correspond to very (infinitely in the zero-order approximation) excited states of the problem **2)** (or conversely). Higher-order corrections do not change this statement. Thus different problems generate different branches of the energy spectrum of the original set of equation. In this respect the following special case differs essentially from the previous ones.

**3).** Let  $r_{2\infty} = r_{1\infty}$  and  $b_{2\infty} = b_{1\infty}$  but  $\rho_2 - \rho_1 = O(\lambda^n)$  and  $\mu_2 - \mu_1 = O(\lambda^n)$ ,  $n \geq 2$ . Then  $w_2 = O(\lambda^2)$ . Both the equations (42) and (43) have similar structure and should be treated on the same footing. The use of common variables  $\xi$ ,  $\epsilon$  defined by (22) and (26) is appropriate to this case. In the zero-order approximation we obtain the coupled pair of wave equations (on the contrary to the cases **1)** and **2)** where we had a single wave equation). In physically meaningful cases (of Section 5, for example) they have the form:

$$\left\{ d^2/d\xi^2 + \kappa\epsilon - \nu_1 - \omega^2\xi^2 \right\} \psi_1(\xi) = \chi\psi_2(\xi), \quad (49)$$

$$\left\{ d^2/d\xi^2 + \kappa\epsilon - \nu_2 - \omega^2\xi^2 \right\} \psi_2(\xi) = \chi\psi_1(\xi), \quad (50)$$

where  $\chi = \lim_{\lambda \rightarrow 0} y = \text{const}$ , and parameters  $\nu_i$ ,  $\kappa$  and  $\omega$  are related to the functions  $w_i$  ( $i = 1, 2$ ) by the equation of the type of (32), (33)

and (34). The equations (49), (50) can be evidently split and reduced to the pair of equations of the type (31) with parameters  $\tilde{\nu}_i = \{\nu_1 + \nu_2 \pm \sqrt{(\nu_1 - \nu_2)^2 + 4\chi^2}\}/2$  ( $i = 1, 2$ ). The eigenvalues  $\epsilon$  corresponding to the first and second equations are split by finite constant  $\tilde{\nu}_1 - \tilde{\nu}_2$ . Thus the corresponding states mix in higher orders of perturbation procedure.

## 5. Application: Regge trajectories of mesons

Here we apply the pseudoperturbative treatment of 2BDE in a meson spectroscopy.

It is known [29] that spectra of heavy mesons are well described by the nonrelativistic potential model with QCD-motivated funnel potential  $u(r) = u_l(r) + u_C(r)$ , where

$$u_C(r) = -\alpha/r, \quad \alpha = 0.27 \quad (51)$$

$$u_l(r) = ar, \quad a = 0.25 \div 0.3 \text{GeV}^2. \quad (52)$$

The Coulomb part (51) of this potential describes a nonrelativistic limit of the vector one-gluon exchange interaction while the linear part (52) is suggested by the area law in the lattice approximation of QCD and has presumably scalar or scalar-vector nature.

The description of light meson spectroscopy needs an application of appropriate relativistic models. Most of them are related to the string theory. From the theoretical viewpoint the most interesting are QCD-motivated relativistic models embracing properties of both the heavy and light mesons. Such models should reflect the scalar-vector structure of interaction and should lead to the funnel-type potential in the nonrelativistic limit.

A natural pretender to the relativistic potential model is the 2BDE with a short-range vector potential and a long-range scalar one. At least three general structures of vector potential are used in the literature,

$$U_v(r) = u_v(r), \quad (53)$$

$$U_v(r) = \{1 - \alpha_1 \cdot \alpha_2\} u_v(r), \quad (54)$$

$$U_v(\mathbf{r}) = \left\{ 1 - \frac{1}{2} \alpha_1 \cdot \alpha_2 \right\} u_v(r) + \frac{1}{2} (\mathbf{n} \cdot \alpha_1) (\mathbf{n} \cdot \alpha_2) r u'_v(r), \quad (55)$$

with  $u_v(r) = u_C(r)$  or another short-range potential; here  $u'(r) = du'(r)/dr$ . The potential (53) is only a static part of vector interaction (see [5]). The relativistic vector field kinematics is taken into account in the potential (54) (see [8, 16]) which, for the Coulomb case, was first proposed by Eddington and Gaunt [27, 28]. In the generalization (55) of the

Breit potential [1] retardation terms have been added [8]. Two different scalar potentials,

$$U_s(r) = \beta_1 \beta_2 u_s(r), \quad (56)$$

$$U_s(r) = \frac{1}{2}(\beta_1 + \beta_2)u_s(r), \quad (57)$$

come from different couplings of scalar mediating field with fermionic fields. The first one (56) arises from the Yukawa interaction (see [6] while the second one (57) corresponds to so called “minimal” coupling [15]. The latter and also two next potentials can be treated as static approximation of various QFT-motivated scalar quasipotentials [17, 30–32]:

$$U_s(r) = \frac{1}{2}(1 + \beta_1 \beta_2)u_s(r), \quad (58)$$

$$U_s(r) = \frac{1}{4}(1 + \beta_1)(1 + \beta_2)u_s(r). \quad (59)$$

The perturbative treatment of Breit-type equations has been used for calculating a fine splitting in spectra of heavy mesons [14, 16]. Light meson spectra are essentially relativistic and need a nonperturbative statement of the problem which is inconsistent because of non-physical singularities of radial equations. To avoid this difficulties in numerical calculations one is forced to invent sophisticated potentials and impose rather artificial boundary conditions [15].

Using the pseudoperturbative treatment of 2BDE with different combinations of potentials (51)–(59) we obtain analytical expressions for meson mass spectra and estimate a role of general structure and input parameters of potentials in the model. We consider mass spectra of lightest mesons (containing  $u$  and  $d$  quarks only) and try to reproduce their following general features:

- i). Mass spectra of light mesons fall into the family of straight lines in the  $(E^2, j)$ -plane known as Regge trajectories.
- ii). Regge trajectories are parallel; slope parameter  $\sigma$  is an universal quantity,  $\sigma = 1.15 \text{ GeV}^2 = (4 \div 4.5)a$ .
- iii). Nonrelativistic classification of light mesons as  $(n^{2s+1}\ell_j)$ -states of quark-antiquark system is adequate; i.e., radial quantum number  $n_r = n - \ell - 1$  enumerates leading ( $n_r = 0$ ) and daughter ( $n_r = 1, 2, \dots$ ) Regge trajectories, spin  $s = 0, 1$  corresponds to mass singlets and triples etc.
- iv). Spectrum is  $ls$ -degenerated, i.e., masses are distinguished by  $\ell$  and  $n_r$  (not by  $j$ ).
- v). States of different  $\ell$  possess an accidental degeneracy which fact causes a tower structure of the spectrum.
- vi). Hyperfine  $ss$ -splitting is relatively small, about  $5 \div 6\%$  of  $\sigma$ .

For this purpose we use the nonrelativistic potential function (51) and (52) in vector and scalar potentials of different spin structure (53)–(59) and calculate pseudoperturbative spectrum in zero-order approximation. Classification of states then is done using singlet-triplet properties of large-large component of wave function (3) in the nonrelativistic limit.

If the vector short-range interaction is ignored and scalar potentials (56)–(59) are used with  $u_s(r) = ar$  the pseudoperturbative mass (i.e., energy) of meson in zero-order approximation has the following form:

$$E_A^2 = ka[\ell + \frac{1}{2} + \eta(n_r + \frac{1}{2})] + \zeta m_+ \sqrt{2a\ell} + \delta_1 m_+^2 - \delta_2 m_1 m_2 + O(1/\sqrt{\ell}), \quad E_0^2 = E_A^2, \quad E_{\pm}^2 = E_A^2 \pm \varkappa a; \quad (60)$$

here  $m_+ = m_1 + m_2$ , and  $k, \eta, \zeta, \delta_1, \delta_2, \varkappa$  are dimensionless constants depending on the potential chosen.

Four families of energy levels  $E_i$  ( $i = A, 0, -, +$ ) form trajectories in the  $(E^2, \ell)$ -plane which are nearly straight. Indeed,  $\zeta = 0 \div 2$  for all the potentials considered and rest masses  $m_a$  ( $a=1,2$ ) of lightest mesons are small compared to the energy scale  $\sqrt{\sigma}$ . Thus the parameter  $\zeta m_+ \sqrt{a}$  determining a curvature of trajectories is small. Parameters  $\delta_1$  and  $\delta_2$  determine a common shift of all the trajectories and are not important for the present discussion. Below we discuss the calculated values of parameters  $k, \varkappa$  and  $\eta$  determining the slope of trajectories and their degeneracy properties.

In the (56) case  $k = 4$  so that the slope  $\sigma = ka$  matches quite well to that of property ii);  $\eta = 2$  causes the accidental degeneracy typical for the harmonic oscillator; but  $\varkappa = 4$  leads to  $j$ -dependence of energy (not  $\ell$ -dependence) so that the  $ls$ -degeneracy is absent.

In the (57) case  $k = 4$  and  $\eta = 2$ , so that the slope and the accidental degeneracy are the same as in the (56) case;  $\varkappa = 4 - 3\sqrt{2} \approx -0.243$  provides an approximate  $ls$ -degeneracy, with accuracy 6%; the splitting is of order of the actual  $ss$ -splitting (see property vi)).

In the (58) case  $k = \varkappa = 3\sqrt{3} \approx 5.196$ ,  $\eta = \sqrt{3} \approx 1.732$ ; none of these values matches well to properties ii)–vi).

In the (59) case  $k = \sqrt{23 - \sqrt{17}}(7 + \sqrt{17})^2/128 \approx 4.2$  provides the best fit of  $\sigma$  to that of property ii);  $\eta = (\sqrt{17} - 3)\sqrt{102 + 26\sqrt{17}}/8 \approx 2.03$  leads to nearly precise oscillator-like degeneracy, with accuracy 1.5%;  $\varkappa = 0$  provides exact  $ls$ -degeneracy.

Taking into account the vector short-range interaction (one of potentials (53)–(55) with  $u_v(r) = u_C(r)$ ) results in a parallel shift of Regge trajectories. The value of the shift is of order  $\alpha a$ , it depends on the vector potential chosen and is different (in general) for different trajectories

$E_i$  ( $i = A, 0, -, +$ ).

It has been proved in the framework of single-particle Dirac equation a possibility of confinement by means of vector and equally mixed vector-scalar long-range interactions [33–35]. We examined these cases in 2BDE approach using different vector potentials (53)–(55) with  $u_v(r) = ar$ . Corresponding zero-order pseudo-perturbative spectra have a form similar to (60). The difference is that  $k = k_i = 8 \div 12$  is as twice or more as desired, and is different for  $i = A, 0, -, +$  (i.e., trajectories are not parallel).

## 6. Summary

The Breit equation and its generalizations (2BDE) possess non-physical singularities. In some cases these points lay far from the physically important domain but they make a boundary problem incorrect or physically improper [16, 17]. In order to avoid this difficulty and to use the 2BDE in the relativistic bound state problem, especially for the case of strong coupling, we develop the  $1/j$  expansion method.

The method is based on the large- $N$  or large- $\ell$  techniques applicable to the radial Schrödinger equation. In our case the 2BDE is reduced to the coupled pair of quasipotential-type equations which structure causes principal modification of known techniques. Other changes are related to the fact that the equations represent a nonlinear spectral problem with cumbersome quasipotentials.

We apply this pseudoperturbative method to the 2BDE with the linear+Coulomb potential of different scalar-vector structure. In all cases in the zero-order approximation it was obtained the Regge trajectories which are linear asymptotically. Linear potentials of two scalar structures (57) and (59) which was discarded in Ref. [17] as nonphysical (because of singularities in 2BDE) reproduce well in our case general properties of light meson spectra. In particular, the slope  $\sigma = ka$  of light meson trajectories fit experimental value well if the parameter  $a$  is taken from the nonrelativistic potential model [29]. Third linear potential of Ref. [17] (with no singularities in 2BDE) does not match to experimental data.

## Acknowledgements

The author would like to thank Professors V. Tretyak, I. Simenog and Dr. Yu. Yaremko for helpful suggestions and critical comments.

## References

1. Breit G., The effect of retardation on the interaction of two electrons, *Physical Review*, 1929, V. 34, N 4, 553–573.
2. Barut A. O. and Komy S., Derivation of nonperturbative relativistic two-body equation from the action principle in quantum electrodynamics, *Fortschritte der Physik*, 1985, V. 33, N 6, 309–318.
3. Barut A. O. and Ünal N., A new approach to bound-state quantum electrodynamics, *Physica*, 1987, V. 142A, N 1–3, 467–487.
4. Grandy W. T. Jr., *Relativistic Quantum Mechanics of Leptons and Fields*, Dordrecht-Boston-London, Kluwer Academic Publishers, 1991.
5. Darewych J. W. and Di Leo L., Two-fermion Dirac-like eigenstates of the Coulomb QED Hamiltonian, *Journal of Physics A*, 1996, V. 29, N 21, 6817–6841.
6. Darewych J. W., Few-particle eigenstates in the Yukawa model, *Condensed Matter Physics*, 1998, V. 1, N 3(15), 593–604.
7. Darewych J. W. and Duviryak A., Exact few-particle eigenstates in partially reduced QED, *Physical Review A*, 2002, V. 66, N 3, 032102-1–032102-20.
8. Duviryak A. and Darewych J. W., Variational wave equations of two fermions interacting via scalar, pseudoscalar, vector, pseudovector and tensor fields, *Central European Journal of Physics*, 2005, V. 3, N 3, 1–17.
9. Fushchich W. I. and Nikitin A. G., On the new constants of motion for two- and three-particle equations, *Journal of Physics A*, 1990, V. 23, N 11, L533–L535.
10. Nikitin A. G. and Fushchich W. I., Non-Lie integrals of the motion for particles of arbitrary spin and for systems of interacting particles, *Teoreticheskaya i Matematicheskaya Fizika*, 1991, V. 88, N 3, 406–515 [English transl.: *Theoretical and Mathematical Physics*, Vol. 88, N 3, 960–967].
11. Simenog I. V. and Turovsky A. I., A relativistic model of the two-nucleon problem with direct interaction, *Ukraïn. Fiz. Zh.*, 2001, V. 46, N 4, 391–401 (in Ukrainian).
12. Simenog I. V. and Turovsky A. I., The model of deuteron in Dirac-Breit approach with direct interaction, *Journal of Physical Studies*, 2004, V. 8, N 1, 23–34 (in Ukrainian).
13. Krolkowski W., Relativistic radial equations for 2 spin-1/2 particles with a static interaction, *Acta Physica Polonica B*, 1976, V. 7, N 7, 485–496.

14. Childers R. W., Effective Hamiltonians for generalized Breit interactions in QCD, *Physical Review D*, 1987, V. 36, N 2, 606–614.
15. Brayshaw D. D., Relativistic description of quarkonium, *Physical Review D*, 1987, V. 36, N 5, 1465–1478.
16. Tsibidis G. D., Quark-Antiquark Bound States and the Breit Equation, LANL e-Print archive, 2000, <http://xxx.lanl.gov/abs/hep-ph/0007143>.
17. Khelashvili A. A., Radial quasipotential equation for a fermion and antifermion and infinitely rising central potentials, *Teoreticheskaya i Matematicheskaya Fizika*, 1982, V. 51, N 2, 201–210 [English transl.: *Theoretical and Mathematical Physics*, 1982, V. 51, N 2, 447–453].
18. Crater H. W., Wong C. W. and Wong C.-Y., Singularity-free Breit equation from constraint two-body Dirac equations, *International Journal of Modern Physics E*, 1996, V. 5, N 4, 589–615.
19. Mlodinov L.D. and Shatz M. P., Solving the Schrödinger equation with use of  $1/N$  perturbation theory, *Journal of Mathematical Physics*, 1984, V. 25, N 4, 943–950.
20. Imbo T., Pagnamenta A. And Sukhatme U., Energy eigenstates of spherically symmetric potentials using the shifted  $1/N$  expansion, *Physical Review D*, 1984, V. 29, N 8, 1669–1681.
21. Vakarchuk I. O., The  $1/N$ -expansion in quantum mechanics. High-order approximations, *Journal of Physical Studies* 2002, V. 6, N 1, 46–54.
22. Mustafa O. and Barakat T., Nonrelativistic shifted- $l$  expansion technique for three- and two-dimensional Schrodinger equation, *Communications in Theoretical Physics*, 1997, V. 28, N 3, 257–264.
23. Mustafa O. and Barakat T., Relativistic shifted- $l$  expansion technique for Dirac and Klein-Gordon equations, *Communications in Theoretical Physics*, 1998, V. 29, N 4, 587–594.
24. Todorov I. T., Quasipotential equation corresponding to the relativistic eiconal approximation, *Physical Review D*, 1971, V. 3, N 10, 2351–2356.
25. Rizov V. A., Sazdian H. and Todorov I. T., On the relativistic quantum mechanics of two interacting spinless particles, *Annals of Physics (NY)*, 1985, V. 165, N 1, 59–97.
26. Duviryak A., Heuristic models of two-fermion relativistic systems with field-type interaction, *Journal of Physics G*, 2002, V. 28, N 11, 2795–2809.
27. Eddington A. S., The charge of an electron, *Proceedings of the Royal Society of London, Series A*, 1929, V. 122, N 789, 358-369.
28. Gaunt J. A., The Triplets of Helium, *Philosophical Transactions of*

- the Royal Society of London. Series A*, 1929, V. 228, 151-196; *Proceedings of the Royal Society of London, Series A*, 1929, V. 122, N 790, 513-532.
29. Lucha W., Schoberl F. F. and Gromes D., Bound states of quarks, *Physics Reports*, 1991, V. 200, Issue 4, 127-240.
30. Salpeter E. E., Mass corrections to the fine structure of hydrogen-like atoms, *Physical Review*, 1952, V. 87, N 2, 328–343.
31. Faustov R. N., The proton structure and hyperfine splitting of hydrogen energy levels, *Nuclear Physics*, 1966, V. 75, N 3, 669–681.
32. Khelashvili A. A., Quasipotential Equation for the System of Two Particles with Spin  $1/2$ , *Communications of the Joint Institute for Nuclear Physics*, P2-4327, Dubna, 1969 (in Russian).
33. Long C. and Robson D., Bound states of a relativistic quark confined by a vector potential, *Physical Review D*, 1983, V. 27, N 3, 644–646.
34. Baric N. and Jena S. N., Lorentz structure vs relativistic consistency of an effective power-law potential model for quark-antiquark systems, *Physical Review D*, 1982, V. 26, N 9, 2420–2429.
35. Haysak I., Lengyel V. et al., Quark masses in the relativistic analytic model, *Ukrain. Fiz. Zh.*, 1996, V. 41, N 3, 370–372 (in Ukrainian).



Препринти Інституту фізики конденсованих систем НАН України розповсюджуються серед наукових та інформаційних установ. Вони також доступні по електронній комп'ютерній мережі на WWW-сервері інституту за адресою <http://www.icmp.lviv.ua/>

The preprints of the Institute for Condensed Matter Physics of the National Academy of Sciences of Ukraine are distributed to scientific and informational institutions. They also are available by computer network from Institute's WWW server (<http://www.icmp.lviv.ua/>)

Аскольд Андрійович Дувіряк

МЕТОД РОЗКЛАДІВ ЗА ВЕЛИКИМ  $j$  ДЛЯ ДВОЧАСТИНКОВОГО  
РІВНЯННЯ ДІРАКА

Роботу отримано 26 грудня 2005 р.

Затверджено до друку Вченою радою ІФКС НАН України

Рекомендовано до друку семінаром відділу теорії металів та сплавів

Виготовлено при ІФКС НАН України

© Усі права застережені